

Cantharone Care

1. Keep the treated areas covered with a bandage or tape if applied at your visit. This will prevent potentially getting the medication on any unwanted skin area or eyes. Also avoid getting the treated areas moist or wet as this may spread the Cantharone to unwanted areas.
2. After having Cantharone applied, you should wash it off gently with soap and water in:
 2 hours 4 hours 6-8 hours
 Leave on overnight Once a blister has formed

If you do not remove the chemical, further blistering will occur.

3. For treatment of the blister sites, cleanse daily with an antibacterial soap and apply Polysporin or Vaseline ointment and a band-aid. There may be some discomfort from the blister sites. Motrin or Tylenol may be used if needed.

At the first treatment, only a few lesions may be treated as a “test” to make sure the patient does not have a severe reaction to Cantharidin (this is rare). If a severe reaction does occur (large blistering, intense redness, swelling and pain), please call the office at 205.871.7332. Another treatment option will have to be used instead.

If the area becomes very red, painful, and drains pus, call our office at 205.871.7332 for further instructions.

Homewood
T: 205.871.7332
F: 205.871.7336
3415 Independence Dr, Suite 200
Birmingham, AL 35209

Greystone
T: 205.678.7518
F: 205.677.2079
5406 US-280, Suite A-100,
Birmingham, AL 35242

info@skinwellness.com
skinwellness.com